

artikel wiryono

Dari: wiryono wiryono (wiryonogood@yahoo.com)

Kepada: biodiversitas@gmail.com; unsjournals@gmail.com

Tanggal: Kamis, 21 Juli 2016 12.05 GMT+7

Dear **Editor-in-Chief,**

I herewith enclosed a research article,

Title:

The survival rate and one-year growth of *Shorea javanica*, *Shorea macrobalanos* and *Hopea mengarawan* in coal mined land in Central Bengkulu, Indonesia

Author(s) name:

Wiryono, Hery Suhartoyo, Ali Munawar

For possibility publication on the journal:

(fill in *Biodiversitas* or *Nusantara Bioscience*)

Biodiversitas

Novelty:

(state your claimed novelty of the findings versus current knowledge)

This article shows that dipterocarps can grow well in mined land. Most mined land reclamation use exotic fast growing species.

Statements:

This manuscript has not been published and is not under consideration for publication to any other journal or any other type of publication (including web hosting) either by me or any of my co-authors.

Author(s) has been read and agree to the Ethical Guidelines.

List of five potential reviewers

(He/she should have Scopus ID and come from different institution with the authors)

Place and date:

Bengkulu, 29 July 2016

Sincerely yours,

(fill in your name, no need scanned autograph)

Wiryono

Address

(Fill in your institution's name and address, your personal cellular phone and email)

The Department of Forestry, Faculty of Agriculture, University of Bengkulu
Jalan WR Supratman Bengkulu 38371A. phone 621367464675
email:wiryonogood@yahoo.com

 111kB

Review report for “The survival and growth of Dipterocarps in coal mined land”

By IT at 10 August 2016

Article type: Short communication

This manuscript reported the survival and growth properties of one-year-old seedlings of three dipterocarps trees. Soil and growing conditions were well documented and I think it is suitable to publishing for Biodiversitas after minor collections. Most of them have already been revised in the attached MS Word file (The survival rate and one-year growth of *Shorea javanica*_IT.doc) by tracking function. The followings are the remaining ones.

All Figures should be appropriately cited in the main text.

Figures 2 and 3 do not need to be drawn by using colors. They should be drawn in black and white.

All references should be cited by an appropriate manner.

1 **Short Communication:**
2 **The survival rate and one-year growth of *Shorea javanica*, *Shorea***
3 ***macrobalanos* and *Hopea mengarawan* in coal mined land in Central**
4 **Bengkulu, Indonesia**

5
6
7
8
9 **Abstract.** Dipterocarp trees used to dominate the lowland forest of Sumatera and Kalimantan. Currently, however, dipterocarp trees are
10 rare due to deforestation of natural forest. One major cause of deforestation in Sumatera and Kalimantan is coal mining. Rehabilitation
11 of coal mined soil is usually done using fast-growing alien species. We tried to restore a piece of mined land in Central Bengkulu, using
12 commercially valuable species of Dipterocarpaceae, namely *Shorea javanica* Koord. and Valet., *Shorea macrobalanos* Ashton and
13 *Hopea mengarawan* Miq. In this article, we presented the survival rate and growth of the one year old seedlings of these three species
14 within one year of observation. Of the three species, *S. macrobalanos* had the highest survival rate (93%), followed by *S. javanica*
15 (80%) and *H. mengarawan* (77%). Within a year, the one year old seedlings of *S. macrobalanos* grew 452% in height, significantly
16 higher than that of *S. javanica* (221%) and of *H. mengarawan* (119%). *Shorea macrobalanos* also had the highest growth in diameter
17 within a year, namely 337%, followed by *S. javanica* (145%) and *H. mengarawan* (135%). -It can be concluded that within a year of
18 observation, the three species of dipterocarp could grow relatively well in mined land. It is therefore recommended that in the restoration
19 of mined land in Sumatera ~~use~~ economically valuable native species of dipterocarp should be used instead of fast-growing alien species.

20 **Key words:** coal mined soil, dipterocarpaceae, restoration

21 **Running title:** The survival and growth of dipterocarps in coal mined land

22 **INTRODUCTION**

23 Trees of the Dipterocarpaceae family used to dominate the lowland forest of Indonesia (Whitmore 1984), but most of
24 them are now threatened with the disappearing of lowland forest in Indonesia. One of industries contributing to the
25 deforestation is mining. Between 2000 and 2010, mining industries caused 0.3 million hectares of forest loss, resulting in
26 3.6- 4.4 Mega ton of carbon emission (Abood et al. 2015). Mining activities not only remove the vegetation, but also
27 drastically destroy the ecosystem. Topsoil was removed and the materials previously buried deep underground were
28 brought to surface. Mine soil, is generally unfavorable for plant growth and must be amended before revegetation
29 (Bradshaw 1997, Lottermoser 2010).

30 The Indonesian government regulations require that the mining companies rehabilitate the mined land in order to
31 restore the ecosystem functions. Usually, exotic fast growing species, such *Acacia mangium* (Suhartoyo et al. 2012),
32 *Paraserianthes falcataria*, *Sesbania grandiflora* (Munawar et al. 2011) are used for the revegetation of mined land. While
33 these species can restore some ecosystem functions of the degraded forest, the establishment of these species changes the
34 species composition of the forest. According to the Society of Ecological Restoration (SER) one of nine ecosystem
35 attributes as the criteria of restoration success is that indigenous species must be present in the area (Clewell and Aronson
36 2007). Several researchers have tried to use local fast growing species for revegetation of mined land in Kalimantan
37 (Adman et al. 2012) and found that some local fast growing species were able to survive and grow in mined land. Those
38 species, however, had low economical value.

39 Considering the lack of study of coal mined land restoration using dipterocarpaceae, we tried to plant three species of
40 dipterocarpaceae in coal mined land in Central Bengkulu District ~~which had been planted with~~ where raintree (*Samanea*
41 *saman*) had already been planted. The three species were *Shorea javanica* Koord. and Valet., *Shorea macrobalanos* Ashton
42 and *Hopea mengarawan* Miq. In some literature *H. mengarawan* is written as *H. mengerawan* Miq. *Shorea javanica* is
43 naturally found in primary and secondary forest in Sumatra and Central Java, but it has been planted in many countries of
44 Southeast Asia (Orwa et al. 2009). *Shorea macrobalanos* used to be endemic to Sarawak and East Kalimantan, and it is
45 categorized as critically endangered species in IUCN Redlist (Ashton 1998a; IUCN 2016-). *Hopea mengarawan* is native
46 to Indonesia and Malaysia (Fern 2016) and it is also categorized as critically endangered in IUCN Redlist (IUCN 2016).
47 The objective of this article was to report the survival rate and the growth of one year old seedlings of the three species
48 within a year of observation.

MATERIALS AND METHODS

Study site

This study was conducted in the mining concession area of PT Inti Bara Perdana in Taba Penanjung Sub-district, Central Bengkulu District, Bengkulu. The site was approximately at 300m altitude, with annual rainfall above 3000 mm. The site was revegetated with raintree (*Samanea saman*) four years earlier. The average canopy opening of the four-year old raintree stand was 40%, so there was some light reaching the forest floor. The light intensity on forest floor under the shade at noon varied between 800 to 3000 lux. The temperature at noon was 28-30°C and the average relative humidity was 82%. During the driest month of 2015, namely October, the raintrees shed most of their leaves, so the canopy was open, and consequently the temperature and light intensity increased while the relative humidity dropped.

The land of this site was not levelled, but consisted of many mounds and depressions. The seedlings were planted on mounds in order to prevent waterlogging and to avoid compacted soil. Seedlings of three species of dipterocarp were planted, namely *Shorea macrobalanos* Ashton, *Shorea javanica* Koord. and Valet. and *Hopea mengarawan* Miq. The seedlings were approximately one year old. At the beginning of planting, a kilogram of compost was given for each planting hole. Afterward, every two months NPK fertilizer was given to each plant. During the months of July-September 2015, the seedlings were watered every two weeks, and during October they were watered every three days.

Every month, the height and diameter of seedlings were measured. Soil chemical and physical properties were analyzed in Soil Laboratory of the Agriculture Faculty, University of Bengkulu.

Data analysis

The data of height growth and diameter growth were analyzed using ANOVA to know whether there were growth differences among species. If there were significant differences, further analyses were done using Least Square Differences (LSD) to know which species had higher growth than the others.

Figure 1. Study site in Taba Penanjung Sub-district, Central Bengkulu District, Bengkulu Province, Indonesia

RESULTS AND DISCUSSION

Shorea macrobalanos had the high survival rate among the three species (Table 1). Only two out of 30 plants died during the first year of planting. *Shorea javanica* and *Hopea mengarawan* had similar survival rate. *Shorea macrobalanos* also had the highest growth of diameter and height within a year of observation, and there was no statistically significant difference between *S.javanica* and *H.mengarawan* in diameter and height growth (Tables 2 and 3). Within a year, *S.macrobalanos* grew 452% in height and 337% in diameter, while *S.javanica* 221% and 245%, and *H.mengarawan* 119% and 135%. In general, the results showed that the three dipterocarp species grew relatively well in the study site.

The success of terrestrial ecosystem restoration is influenced by the plant-soil interactions (Eviner and Hawkes 2008). Plant growth in mined land is sometimes constrained by soil compaction (Sheoran et al. 2010). Soil compaction may impede root penetration, reduce aeration, slow down movement of nutrients and water, and cause the buildup of toxic gases in the rhizosphere (Brady and Weil. 2002). In the United States, the passing of Surface Mining Control and Reclamation Act, which require the mined land to be returned to its original contour, had resulted in soil compaction due to the use of heavy equipment. As a result, the reclaimed mined land was mostly grown with grasses and herbaceous vegetation, while tree establishment was impeded by the high soil density (Fields-Johnson et al. 2014). To improve the establishment of trees, the soil compaction must be reduced, for example through ripping (Ashby 1997, Fields-Johnson et al. 2014). This problem of soil compaction, however, was not found in our study site, because the mining company did not level the soil. The dipterocarp seedlings were planted in piles of soil which were not compacted. The bulk density of the soil samples ranged between 1.49 and 1.60 g cm⁻³ and the soil particle density was 1.84-1.88 g cm⁻³ (Table 4). The bulk density in this study site was considered normal, because the mineral soil may have bulk density of 1.0-2.0 g cm⁻³ (Chapin et al 2011) and the soil particle density was lower than the average particle density of mineral soil, which is 2.5-2.7 g cm⁻³. With normal soil density, the roots of dipterocarp seedlings would not have problem in penetrating the soil. Also, because the seedlings were planted on the mounds, there was no problem of water saturation during rainy day.

The chemical properties of soil in mined land is in general unfavorable for plant growth. Mine soil which sometimes contains high concentration of sulfur can have a pH of 2.3 – 3.5 which lead to higher availability of toxic metal (Sheoran et al 2010). In our study site, the chemical soil was very acidic with a pH of 4.0- 4.2 (Table 5), but the pH was not extremely low. Even in primary tropical rain forest, the soil is usually also acidic. In SouthWest China, Li et al (2012) found the pH of primary tropical rain was 4.53.

In mine soil, the nutrient availability is usually low, and so it was in our study site (Table 5). However, at the beginning of planting, manure was given for each seedling and subsequently, every two months, NPK fertilizer at the amount of 15 gram was given for each seedling. So, the nutrient content of the soil in the rhizosphere of the dipterocarp seedlings was

certainly higher than that in the surrounding. Soil amendment could increase nutrient availability for plants in mine soil (Asensio et al 2014).

Beside soil properties, light intensity also influences the establishment of plants in reclaimed mined land. Every species requires different need of light intensity for germination and establishment. Ashton (1998b) said that dipterocarp species could be broadly classified as shade-intolerant, but some species were shade-tolerant. Generally, the seeds require partial shade for germination and early survival, but the seedlings require an increase in light for establishment and growth. Bawalsyah et al (2015) in their study found in West Sumatra that there was no difference in the growth of *H.mengarawan* in different light intensity, which was 100% and 40%. In our study, the canopy of raintree was not fully closed. The light intensity on the forest floor, under the shade at noon varied between 800-3500 lux.

The plant growth and survival are also influenced by its interaction with other organisms. Attack from pathogen to the dipterocarp seedlings was not found in our study, while minor herbivory was found but not lethal. Some plants died because they were uprooted by wild pigs. Fortunately, the leaves of the three dipterocarps were not palatable to large herbivore, so depredation from large herbivores was absent.

The most important threat came from competition with weeds, such as *Mikania micrantha*, *Synedrella nodiflora*, *Chromolaena odorata*, *Nephrolepis biserrata* and *Mimosa pudica*. These weeds can easily overgrow and enclose the dipterocarps seedlings, depriving the seedlings of the light. The worst weed was *Mikania micrantha*. Native to America (Holm et al 1977), this twinning perennial weed would twin its stem around the seedlings' stem and branches. To ensure the establishment of the dipterocarp, every two weeks we manually removed the weeds surrounding the dipterocarps. In the US, Ashby (1997) found that the use of herbicide helped improve tree establishment in mined land.

It can be concluded that with proper care, *Shorea javanica*, *Shorea macrobalanos* and *Hopea mengarawan* could grow relatively well in coal mined under the rain tree stand in Central Bengkulu. It is recommended that dipterocarp trees be used to revegetate the mined land in Sumatra.

Formatted: Font: Italic

Formatted: Font: Italic

ACKNOWLEDGEMENTS

We gratefully acknowledge the help of several parties for this study. This study was funded by the Directorate General of Higher Education. The management of the mining company, Inti Bara Perdana Ltd, permitted us to do this study in their mining concession area. M. Fajrin Hidayat made the map of the Syudy site for this article.

REFERENCES

- Abood SA, Lee JSH, Burivalova Z, Garcia-Ulloa J, Koh LP. 2015. Relative contribution of the logging, fiber, oil palm and mining industries to forest loss in Indonesia. *Conservation letters* 8(1):58-67.
- Adman B, Hendrarto B, Sasongko DP. 2012. Pemanfaatan jenis lokal cepat tumbuh untuk pemulihan lahan pascatambang batubara. (Studi kasus di PT Singlurus Pratama, Kalimantan Timur). *Jurnal Ilmu Lingkungan* 10(1): 19-25.
- Asensio V, Vega FA, Covelo EF. 2014. Changes in the Phytoavailability of Nutrients in Mine Soils after Planting Trees and Amending with Wastes V. *Water Air Soil Pollut* 225:1995
- Ashby WC (1997) Soil ripping and herbicides enhance tree and shrub restoration on stripmines. *Restor Ecol* 5:169–177
- Ashton P. 1998a. *Shorea macrobalanos*. *The IUCN Red List of Threatened Species 1998*: e.T31921A9668585. <http://dx.doi.org/10.2305/IUCN.UK.1998.RLTS.T31921A9668585.en>
- Ashton P. 1998b. Seedling ecology of mixed dipterocarp forest. In: Appanah S, Turnbull JM. (Eds.). *A Review of Dipterocarps. Taxonomy, ecology and silviculture*. CIFOR.
- Bawalsyah E, Suwirman, Noli ZA. 2015. The Growth of *Hopea mengarawan* Miq. Seedling Under Different Light Intensity. *Jurnal Biologi Universitas Andalas (J. Bio. UA.)* 4(2): 90-95
- Bradshaw AD. 1997. What we mean by restoration. In Urbanska KM, Webb NR, Edwards PJ. *Restoration Ecology and Sustainable Development*. Cambridge University Press.
- Brady NC, Weil RR. 2002. The nature and properties of soil. Thirteenth ed. Pearson Education, Upper Saddle River, NJ.
- Chapin FS, Matson PA, Vitousek PM. 2011. Principles of terrestrial ecosystem ecology. Springer. New York.
- Clewell AF, Aronson J. 2007. *Ecological Restoration. Principles, Values and Structure of an Emerging Profession*. Island Press. Washington, D.C.
- Eviner VT, Hawkes CV. 2008. Embracing Variability in the Application of Plant–Soil Interactions to the Restoration of Communities and Ecosystems. *Restoration Ecology* Vol. 16 (4): 713–729
- Fern K. 2016. Useful tropical plants. *Hopea mengarawan*. <http://tropical.ferns.info/viewtropical.php?id=Hopea+mengarawan>. Downloaded on July 11, 2016.
- Fields-Johnson CW, Burger JA, Evans DM, Zipper CE. 2014. Ripping Improves Tree Survival and Growth on Unused Reclaimed Mined Lands. *Environmental Management* 53:1059–1065
- Holm LG, Plucknett DL, Pancho JV, Herberger J.P. 1977. *The World's worst weeds*. The University Press of Hawaii. Honolulu.

- 159 IUCN 2016. The IUCN Red List of Threatened Species. Version 2016-1. <www.iucnredlist.org>. Downloaded on **11 July**
 160 **2016**
- 161 Li H, Ma Y, Liu W. Liu W. 2012 Soil Changes Induced by Rubber and Tea Plantation Establishment: comparison with
 162 Tropical Rain Forest Soil in Xishuangbanna, SW China. Environmental Management 50:837–848
- 163 Lottermoser BG. 2010. Mine Wastes. Characterization, treatment and environmental impacts. 3rd Ed. Springer-Verlag.
 164 Berlin.
- 165 Munawar A, Indramawan, Suhartoyo H (2011) Litter production and decomposition rate in the Reclaimed Mined Land
 166 under Albizia and Sesbania stands and their effects on some soil chemical properties. Journal of Tropical Soils (16) 1:
 167 1-6
- 168 Orwa C, A Mutua, Kindt R , Jamnadass R, S Anthony. 2009 Agroforestry Database:a tree reference and selection guide
 169 version 4.0 (<http://www.worldagroforestry.org/sites/treedbs/treedatabases.asp>)
- 170 Sheoran V, Sheoran AS, Poonia P. 2010. Soil reclamation of abandoned mined land by revegetation: a review.
 171 International Journal of Soil, Sediment and Water: Vol. 3: Iss. 2, Article 13.
- 172 Suhartoyo H, Munawar A, Wiryo. 2012. Returning biodiversity of rehabilitated forest on a coal mined site at Tanjung
 173 Enim, South Sumatra. Biodiversitas vol.13 (1):13-17.
- 174 Whitmore TC. 1984. Tropical Forest of the Far East. Clarendon Press. Oxford.

175
 176 Table 1. The survival rate of three Dipterocarp species in coal mined land during a year of observation
 177

Species	No of plants alive	Survival rate (%)
<i>Shorea macrobalanos</i>	28	93
<i>Shorea javanica</i>	24	80
<i>Hopea mengarawan</i>	23	77

178
 179 The growth of height of the three species in
 180
 181
 182

183 Figure 2. The plant height of three Dipterocarp species in coal mined land during a year of observation
 184
 185
 186
 187
 188

189 Figure 3. The plant diameter (in mm) of three Dipterocarp species in coal mined land during a year of observation
 190

191 Table 2. The growth of height (%) of three Dipterocarp species in coal mined land during a year of observation
 192
 193

Species	Height increase in a year (%)
<i>Shorea macrobalanos</i>	452a
<i>Shorea javanica</i>	221b
<i>Hopea mengarawan</i>	119b

194 Note: numbers followed by the same letter are not significantly different at p of 5% level.
 195
 196

197 Table 3. The growth of diameter (%) of three Dipterocarp species in coal mined during a year of observation.
 198
 199

Species	Diameter increase in a year (%)
<i>Shorea macroblanos</i>	337a
<i>Shorea javanica</i>	145b
<i>Hopea mengarawan</i>	135b

200 Note: numbers followed by the same letter are not significantly different at p of 5% level.
 201

202 Table 4. Physical properties of soil samples of coal mined land
 203
 204

	Particle density g cm ⁻³	Bulk density	Texture (hygrometer)			
			Sand (%)	Silt (%)	Clay (%)	Class
Sample 1	1.84	1.49	45.51	24.36	30.13	Sandy clay loam-clay loam
Sample 2	1.88	1.60	76.24	8.68	15.08	Sandy loam

205 Table 5. The chemical properties of soil samples of mined land

205
206

Soil samples	pH (H ₂ O)	C	N	P ₂ O ₅	K	Ca	Mg	CEC
		%		mg.kg ⁻¹	cmol kg ⁻¹			
Sample 1	4.0 very acidic	1.22 Low	0.05 very low	Un-detected	0.35 Medium	3.34 Low	5.02 Low	15.44 low
Sample 2	4.2 very acidic	1.52 low	0.07 very low	Un-detected	0.29 Medium	2.14 low	5.13 low	14.92 low

207
208

Figure 4a. Hopea mengarawan, 1 month old	Figure 4b. Hopea mengarawan, 11 months old.
--	---

209
210

Re: edited shorea article

Dari: Managing Editor (unsjournals@gmail.com)

Kepada: wiryonogood@yahoo.com

Tanggal: Selasa, 6 September 2016 17.53 GMT+7

Thank you,
Best Regards,

Ahmad Dwi Setyawan

Managing Editor,
- Biodiversitas, Journal of Biological Diversity (biodiversitas.mipa.uns.ac.id) (SCOPUS, DOAJ)
- Nusantara Bioscience (biosains.mipa.uns.ac.id/nusbioscience.htm) (Web of Science (ESCI), DOAJ)

Chairman/Co-Chairman

- National Seminar & International Conference on Biodiversity, <http://biodiversitas.mipa.uns.ac.id/snmbi.html>

Department of Biology,
Faculty of Mathematics and Natural Sciences, Sebelas Maret University,
Jl. Ir. Sutami 36A Solo 57126, Central Java, Indonesia,
Tel. & Fax. +62-271-663375,
e-mail: unsjournals@gmail.com

2016-09-06 15:15 GMT+07:00 wiryono wiryono <wiryonogood@yahoo.com>:

Dear Pak Ahmad: Please find attached the edited article of shorea. I have corrected some typos (misspelling).

Thank you for publishing this article.

Best Wishes
wiryono

Re: revised article

Dari: Managing Editor (unsjournals@gmail.com)

Kepada: wiryonogood@yahoo.com

Tanggal: Senin, 29 Agustus 2016 21.36 GMT+7

Pls, send us your final ms with tables and figures.

Thank you,
Best Regards,

Ahmad Dwi Setyawan

Managing Editor,
- Biodiversitas, Journal of Biological Diversity (biodiversitas.mipa.uns.ac.id) (SCOPUS, DOAJ)
- Nusantara Bioscience (biosains.mipa.uns.ac.id/nusbioscience.htm) (Web of Science (ESCI), DOAJ)

Chairman/Co-Chairman

- National Seminar & International Conference on Biodiversity, <http://biodiversitas.mipa.uns.ac.id/snmbi.html>

Department of Biology,
Faculty of Mathematics and Natural Sciences, Sebelas Maret University,
Jl. Ir. Sutami 36A Solo 57126, Central Java, Indonesia,
Tel. & Fax. +62-271-663375,
e-mail: unsjournals@gmail.com

2016-08-29 19:29 GMT+07:00 wiryono wiryono <wiryonogood@yahoo.com>:

Pak Ahmad, please find attached our article which has been revised according to the reviewer's suggestion, except for figures, which are not made in black and white. You may change the figures into black and white, if you think it is necessary.

The tables and figures are not included in this article. You may take them from the original article.

Thank you for publishing this article in Biodiversitas.

Sincerely

wiryono

Re: final-revised article

Dari: Managing Editor (unsjournals@gmail.com)

Kepada: wiryonogood@yahoo.com

Tanggal: Selasa, 30 Agustus 2016 07.34 GMT+7

Thank you,
Best Regards,

Ahmad Dwi Setyawan

Managing Editor,
- Biodiversitas, Journal of Biological Diversity (biodiversitas.mipa.uns.ac.id) (SCOPUS, DOAJ)
- Nusantara Bioscience (biosains.mipa.uns.ac.id/nusbioscience.htm) (Web of Science (ESCI), DOAJ)

Chairman/Co-Chairman
- National Seminar & International Conference on Biodiversity, <http://biodiversitas.mipa.uns.ac.id/snmbi.html>

Department of Biology,
Faculty of Mathematics and Natural Sciences, Sebelas Maret University,
Jl. Ir. Sutami 36A Solo 57126, Central Java, Indonesia,
Tel. & Fax. +62-271-663375,
e-mail: unsjournals@gmail.com

2016-08-30 2:36 GMT+07:00 wiryono wiryono <wiryonogood@yahoo.com>:

Mr. Ahmad: please find attached the revised article with tables and figures.

Thanks
wiryono